

## Theorems and Postulates

### Postulate 2-A Protractor Postulate

Given  $\overline{AB}$  and a number  $r$  between 0 and 180, there is exactly one ray with endpoint  $A$ , extending on either side of  $\overline{AB}$ , such that the measure of the angle formed is  $r$ .

### Definition of Right, Acute and Obtuse Angles

$\angle A$  is a right angle if  $m\angle A$  is 90.

$\angle A$  is an acute angle if  $m\angle A$  is less than 90.

$\angle A$  is an obtuse angle if  $m\angle A$  is greater than 90 and less than 180.

### Postulate 2-B Angle Addition

If  $R$  is in the interior of  $\angle PQS$ , then  $m\angle PQR + m\angle RQS = m\angle PQS$ .

If  $m\angle PQR + m\angle RQS = m\angle PQS$ , then  $R$  is in the interior of  $\angle PQS$ .

Vertical angles are congruent.


The sum of the measures of the angles in a linear pair is  $180^\circ$ .

The sum of the measures of complementary angles is  $90^\circ$ .

### Postulate 3-A Ruler

Two points on a line can be paired with real numbers so that, given any two points  $R$  and  $S$  on the line,  $R$  corresponds to zero, and  $S$  corresponds to a positive number.

Point  $R$  could be paired with 0, and  $S$  could be paired with 10.



### Postulate 3-B Segment Addition

If  $N$  is between  $M$  and  $P$ , then  $MN + NP = MP$ .

Conversely, if  $MN + NP = MP$ , then  $N$  is between  $M$  and  $P$ .

**Theorem 4-A  
Pythagorean  
Theorem**

In a right triangle, the sum of the squares of the measures of the legs equals the square of the measure of the hypotenuse.

**Distance Formula**

The distance  $d$  between any two points with coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$  is given by the formula

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}.$$

**Midpoint  
Definition**

The midpoint, **M**, of  $\overline{AB}$  is the point between **A** and **B** such that **AM** = **MB**.

**Midpoint Formula  
Number Line**

With endpoints of A and B on a number line, the midpoint of  $\overline{AB}$  is  $\frac{A+B}{2}$ .

**Midpoint Formula  
Coordinate Plane**

In the coordinate plane, the coordinates of the midpoint of a segment whose endpoints have coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$  are  $(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2})$ .

**Theorem 4-B  
Midpoint Theorem**

If M is the midpoint of  $\overline{PQ}$ , then  $\overline{PM} \cong \overline{MQ}$ .

**Postulate 5-A  
Law of  
Detachment**

If  $p \Rightarrow q$  is true, and  $p$  is true, then  $q$  is true.

**Postulate 5-B  
Law of Syllogism**

If  $p \Rightarrow q$  is true and  $q \Rightarrow r$  is true, then  $p \Rightarrow r$  is true.