# **PROPERTIES OF TRIANGLES**

In this unit, you will begin with a review of triangles and their properties. You will look at classifying triangles by both angles and sides. You will examine the Angle Sum Theorem and other theorems that apply to triangles. In the final part of this unit, you will explore congruency between triangles by examining their corresponding angles and sides and learn a very useful definition of congruent triangles (CPCTC).

Classifying Triangles by Angles

Classifying Triangles by Sides

Angles of a Triangle

**Congruent Triangles** 

## **Classifying Triangles by Angles**


triangle – A triangle is a three-sided polygon.

**right triangle** – A right triangle is a triangle that has one right angle.


**obtuse triangle** – An obtuse triangle is a triangle that has one obtuse angle.


**acute triangle** – An acute triangle is a triangle in which all the angles are acute.

**equiangular triangle** – An equiangular triangle is a triangle in which all three angles are equal in measure.










All are acute angles (angles that measure less than  $90^{\circ}$ ).




All angles measure the same  $(60^\circ)$ .

# Classifying Triangles by Sides


**scalene triangle** – A scalene triangle is a triangle that has all sides measuring different lengths.

**isosceles triangle** – An isosceles triangle is a triangle that has two congruent sides.

equilateral triangle – An equilateral triangle is a triangle that has all sides congruent.



Let's examine a triangle in the coordinate plane to determine what type of triangle it is.



Given  $\Box$  *KLM* . Use the distance formula to determine the type of triangle based on the length of it sides.

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$


$KL = \sqrt{(-5-4)^2 + (4-2)^2}$	$LM = \sqrt{[4 - (-4)]^2 + [2 - (-2)]^2}$	$KM = \sqrt{\left[-5 - (-4)\right]^2 + \left[4 - (-2)\right]^2}$
$d = \sqrt{(-9)^2 + (2)^2}$	$d = \sqrt{(8)^2 + (4)^2}$	$d = \sqrt{(-1)^2 + (6)^2}$
$d = \sqrt{85}$	$d = \sqrt{80}$	$d = \sqrt{37}$

□ *KLM* is a scalene triangle because each side measures a different length.


### Angles of a Triangle

**exterior angle** – An exterior angle is an angle in the exterior of a triangle and is formed with one side of a triangle and an extension of the adjacent side.


**remote interior angles** – Remote interior angles are the interior angles of a triangle that are not adjacent to an exterior angle.



Activity: Look at the angles in the triangle below. Draw a triangle similar to the one shown below. It does not have to be the exact same size. Cut the angles away from the triangle.



Place the angles side by side at one vertex point, so that there are no gaps between the angles as shown below.



The three angles of the triangle total 180°.

*Example 1*: In  $\Box ABC$ , what is the  $m \angle A$  if the  $m \angle C = 95^{\circ}$  and the  $m \angle B = 40^{\circ}$ ?


 $m \angle A + m \angle B + m \angle C = 180$  $m \angle A + 40 + 95 = 180$  $m \angle A + 135 = 180$  $m \angle A = 45$ 

Angle Sum Theorem (Theorem 12-A) Substitution Simplify Subtraction

Theorem 12-B Third Angle Theorem If two of the angles in one triangle are congruent to two of the angles in a second triangle, then the third angles of each triangle are congruent.

Theorem 12-C Exterior Angle Theorem

In a triangle, the measure of an exterior angle is equal to the sum of the measures of the two remote interior angles. Use the figure below to solve the following examples.



*Example 2*: Find the  $m \angle 1$ .

In this problem, we will apply the Exterior Angle Theorem since  $\angle 1$  is the exterior angle of remote interior angles,  $\angle BCE$  and  $\angle E$ , and the measures of these angles are given.

$m \angle BCE = 48^\circ; m \angle E = 87^\circ$	Given
$m \angle 1 = m \angle BCE + m \angle E$	Exterior Angle Theorem (Theorem 12-C)
$m \angle 1 = 48^\circ + 87^\circ$	Substitution
$m \angle 1 = 135^{\circ}$	Simplify

*Example 3*: Find the  $m \angle 2$ .

 $\angle 1$  and  $\angle 2$  are a linear pair because they form a straight line.

$m \angle 2 + m \angle 1 = 180^{\circ}$	Linear pairs are supplementary. (Theorem 10-B)
$m \angle 1 = 135^{\circ}$	Determined in the previous example.
$m \angle 2 + 135^{\circ} = 180^{\circ}$	Substitution
$m \angle 2 = 45^{\circ}$	Subtraction

*Example 4*: Find the  $m \angle 3$ .

 $\angle 2$  and  $\angle 3$  are vertical angles.

$\angle 2 \cong \angle 3$	Vertical angles are congruent. (Theorem 10-H)
$m \angle 2 = m \angle 3$	Definition of congruency.
$m \angle 2 = 45^{\circ}$	Determined in the previous example.
$m \angle 3 = 45^{\circ}$	Substitution ( $m \angle 2 = 45^{\circ}$ )

#### *Example 5*: Find the $m \angle 4$ .

For this problem, we will apply the Third Angle Theorem (12-B) even though we could also use the Triangle Sum Theorem (12-A).

$m \angle D = m \angle E$	Given
$m \angle 2 = m \angle 3$	Vertical angles have same measure.
$m \angle 4 = m \angle C = 48^{\circ}$	Third Angle Theorem (Theorem 12-B)
	(The measure of the third angle of $\Box$ BCE = 48°.)

#### *Example 6*: Find the $m \angle 5$ .

 $\angle 5$  and  $\angle BCE$  are a linear pair because they form a straight line.

$m \angle 5 + m \angle BCE = 180^{\circ}$	Linear pairs are supplementary. (Theorem 10-B)
$m \angle BCE = 48^{\circ}$	Given
$m \angle 5 + 48^\circ = 180^\circ$	Substitution
$m \angle 5 = 132^{\circ}$	Subtraction

**corollary** – A corollary is a theorem that can be easily proved by means of a closely related theorem.

We will now look at some theorems that are closely related to the Angle Sum Theorem (12-A); thus, instead of referring to them as theorems, we will name them corollaries.


Corollary 12-A-1

The acute angles of a right triangle are complementary.

# Corollary 12-A-2

There can be at most one right angle in triangle.

Given: $\Box JKL$  $\angle K$  is a right angleProve: $\angle J$  and  $\angle L$  are acute angles



### Statements

#### Reasons

$\angle K$ is a right angle.	Given
$m \angle K = 90$	Definition of a right angle.
$m \angle K + m \angle L + m \angle J = 180$	Angle Sum Theorem (Theorem 12-A)
$90 + m \angle L + m \angle J = 180$	Substitution Property
$m \angle L + m \angle J = 90$	Subtraction Property
$m \angle L = 90 - m \angle J$	Subtraction Property
$m \angle J = 90 - m \angle L$	Subtraction Property
$\angle$ L and $\angle$ J are acute angles.	Definition of acute angles

Corollary 12-A-3	There can be at most one obtuse angle in triangle.
Corollary 12-A-4	The measure of each angle in an equiangular triangle is 60.

### **Congruent Triangles**


We will now examine congruent triangles through transformations. The first transformation is a slide.

### Slide

 $\Box ABC \cong \Box XYZ$ 

Slide  $\Box ABC$  to the right and down until it fits exactly on top of  $\Box XYZ$ . These two triangles are indeed congruent.

When two geometric figures are congruent, they have congruent corresponding parts.



### Definition of Congruent Triangles (CPCTC)


Two triangles are congruent if and only if their corresponding parts are congruent.

This definition is one that you will use many times in proofs about triangles. Be sure to memorize it!

Corresponding Parts of Congruent Triangles are Congruent!

## Flip (Reflection)

These figures show congruent triangles that are a reflection of each other.




### \*Note: In this figure, $\overline{MK}$ corresponds to itself.

Postulate 12-A

Any segment or angle is congruent to itself. (Reflexive Property)


### Rotation

These figures show congruent triangles that are rotated. The second figure is a 90 degree rotation of the first figure.



In this figure,  $\Box ACD$  is a 180 degree rotation of  $\Box ACB$  and vice versa.

 $\Box ABC \cong \Box ADC$ 



*Example 1*: Complete each correspondence and congruence statement.

$\angle BAC \leftrightarrow$	$\angle BAC \cong$
$\angle B  \leftrightarrow$	$\angle B \cong \_\_\_$
$\angle BCA \leftrightarrow$	$\angle BCA \cong$
$\overline{AB} \leftrightarrow \_\_\_$ $\overline{BC} \leftrightarrow \_\_\_$	$\overline{AB} \cong \underline{\qquad}$ $\overline{BC} \cong \underline{\qquad}$
$\overline{AC} \leftrightarrow$	$\overline{AC} \cong$

Solution:

$\angle BAC \leftrightarrow \angle ACD$	$\angle BAC \cong \angle ACD$	$\overline{AB} \leftrightarrow \overline{CD}$	$\overline{AB} \cong \overline{CD}$
$\angle B  \leftrightarrow \angle D$	$\angle B \cong \angle D$	$\overline{BC} \leftrightarrow \overline{DA}$	$\overline{BC} \cong \overline{DA}$
$\angle BCA \leftrightarrow \angle DAC$	$\angle BCA \cong \angle DAC$	$\overline{AC} \leftrightarrow \overline{AC}$	$\overline{AC} \cong \overline{AC}$

*Example 2*: Which congruence statement in the previous example is an example of Postulate 12-A? Please explain.

Solution:  $\overline{AC} \cong \overline{AC}$ 

Explanation: In the diagram,  $\overline{AC}$  is a part of both triangles; thus, it is congruent to itself. This is an example of the "reflexive" property of congruent segments, Postulate 12-A.