

## Theorems and Postulates

### Postulate 2-A Protractor Postulate

Given  $\overline{AB}$  and a number  $r$  between 0 and 180, there is exactly one ray with endpoint  $A$ , extending on either side of  $\overline{AB}$ , such that the measure of the angle formed is  $r$ .

### Definition of Right, Acute and Obtuse Angles

$\angle A$  is a right angle if  $m\angle A$  is 90.

$\angle A$  is an acute angle if  $m\angle A$  is less than 90.

$\angle A$  is an obtuse angle if  $m\angle A$  is greater than 90 and less than 180.

### Postulate 2-B Angle Addition

If  $R$  is in the interior of  $\angle PQS$ , then  $m\angle PQR + m\angle RQS = m\angle PQS$ .

If  $m\angle PQR + m\angle RQS = m\angle PQS$ , then  $R$  is in the interior of  $\angle PQS$ .

Vertical angles are congruent.


The sum of the measures of the angles in a linear pair is  $180^\circ$ .

The sum of the measures of complementary angles is  $90^\circ$ .

### Postulate 3-A Ruler

Two points on a line can be paired with real numbers so that, given any two points  $R$  and  $S$  on the line,  $R$  corresponds to zero, and  $S$  corresponds to a positive number.

Point  $R$  could be paired with 0, and  $S$  could be paired with 10.



### Postulate 3-B Segment Addition

If  $N$  is between  $M$  and  $P$ , then  $MN + NP = MP$ .

Conversely, if  $MN + NP = MP$ , then  $N$  is between  $M$  and  $P$ .

**Theorem 4-A  
Pythagorean  
Theorem**

In a right triangle, the sum of the squares of the measures of the legs equals the square of the measure of the hypotenuse.

**Distance Formula**

The distance  $d$  between any two points with coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$  is given by the formula

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}.$$

**Midpoint  
Definition**

The midpoint,  $M$ , of  $\overline{AB}$  is the point between  $A$  and  $B$  such that  $AM = MB$ .

**Midpoint Formula  
Number Line**

With endpoints of  $A$  and  $B$  on a number line, the midpoint of  $\overline{AB}$  is  $\frac{A+B}{2}$ .

**Midpoint Formula  
Coordinate Plane**

In the coordinate plane, the coordinates of the midpoint of a segment whose endpoints have coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$  are  $(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2})$ .

**Theorem 4-B  
Midpoint Theorem**

If  $M$  is the midpoint of  $\overline{PQ}$ , then  $\overline{PM} \cong \overline{MQ}$ .

**Postulate 5-A  
Law of  
Detachment**

If  $p \Rightarrow q$  is true, and  $p$  is true, then  $q$  is true.

**Postulate 5-B  
Law of Syllogism**

If  $p \Rightarrow q$  is true and  $q \Rightarrow r$  is true, then  $p \Rightarrow r$  is true.

**Postulate 6-A  
Reflexive  
Property**

Any segment or angle is congruent to itself.

$$\overline{QS} \cong \overline{QS}$$

**Postulate 6-B  
Symmetric  
Property**

If  $\overline{AB} \cong \overline{CD}$ , then  $\overline{CD} \cong \overline{AB}$ .  
If  $\angle CAB \cong \angle DOE$ , then  $\angle DOE \cong \angle CAB$ .

**Theorem 6-A  
Transitive  
Property**

If any segments or angles are congruent to the same angle, then they are congruent to each other.

If  $\overline{AB} \cong \overline{CD}$  and  $\overline{CD} \cong \overline{EF}$ , then  $\overline{AB} \cong \overline{EF}$ .  
If  $\angle 1 \cong \angle 2$  and  $\angle 2 \cong \angle 3$ , then  $\angle 1 \cong \angle 3$ .

**Theorem 6-B  
Transitive  
Property**

If any segments or angles are congruent to each other, then they are congruent to the same angle. (This statement is the converse of Theorem 6-A.)

**Theorem 7-A  
Addition  
Property**

If a segment is added to two congruent segments, then the sums are congruent.

**Theorem 7-B  
Addition  
Property**

If an angle is added to two congruent angles, then the sums are congruent.

**Theorem 7-C  
Addition  
Property**

If congruent segments are added to congruent segments, then the sums are congruent.

**Theorem 7-D  
Addition  
Property**

If congruent angles are added to congruent angles, then the sums are congruent.

**Theorem 7-E  
Subtraction  
Property**

If a segment is subtracted from congruent segments, then the differences are congruent.

**Theorem 7-F  
Subtraction  
Property**

**If an angle is subtracted from congruent angles, then the differences are congruent.**

**Theorem 7-G  
Subtraction  
Property**

**If congruent segments are subtracted from congruent segments, then the differences are congruent.**

**Theorem 7-H  
Subtraction  
Property**

**If congruent angles are subtracted from congruent angles, then the differences are congruent.**

**Theorem 7-I  
Multiplication  
Property**

**If segments are congruent, then their like multiples are congruent.**

**Theorem 7-J  
Multiplication  
Property**

**If angles are congruent, then their like multiples are congruent.**

**Theorem 7-K  
Division  
Property**

**If segments are congruent, then their like divisions are congruent.**

**Theorem 7-L  
Division  
Property**

**If angles are congruent, then their like divisions are congruent.**

**Theorem 10-A**

**Congruence of angles is reflexive, symmetric, and transitive.**

**Theorem 10-B**

**If two angles form a linear pair, then they are supplementary angles.**

**Theorem 10-C**

**Angles supplementary to the same angle are congruent.**

**Theorem 10-D**

**Angles supplementary to congruent angles are congruent.**

**Theorem 10-E**

**Angles complementary to the same angle are congruent.**

**Theorem 10-F**

**Angles complementary to congruent angles are congruent.**

**Theorem 10-G**

**Right angles are congruent.**

**Theorem 10-H**

**Vertical angles are congruent.**

**Theorem 10-I**

**Perpendicular lines intersect to form right angles.**

**Postulate 10-A**

**If two parallel lines are cut by a transversal, then each pair of corresponding angles is congruent.**

**Theorem 10-J**

**If two parallel lines are cut by a transversal, then each of the pair of alternate interior angles is congruent.**

**Theorem 10-K**

If two parallel lines are cut by a transversal, then each pair of alternate exterior angles is congruent.

**Theorem 10-L**

If two parallel lines are cut by a transversal, then each pair of consecutive interior angles is supplementary.

**Theorem 10-M**

If two parallel lines are cut by a transversal that is perpendicular to one of the parallel lines, then the transversal is perpendicular to the other parallel line.

The definition of slope states that, given two points  $(x_1, y_1)$  and  $(x_2, y_2)$ , the slope of a line containing the points can be determined using this formula:

$$m = \frac{y_2 - y_1}{x_2 - x_1} \text{ when } x_2 - x_1 \neq 0$$

**Postulate 11-A**

Two non-vertical lines have the same slope if and only if they are parallel.

**Postulate 11-B**

Two non-vertical lines are perpendicular if and only if the product of their slopes is  $-1$ .

**Postulate 11-C**

If two lines in a plane are cut by a transversal and the corresponding angles are congruent, then the lines are parallel.

**Postulate 11-D**

If there is a line and a point that is not on the line, then there exists exactly one line that passes through the point that is parallel to the given line.

**Theorem 11-A**

If two lines in a plane are cut by a transversal and the alternate interior angles are congruent, then the lines are parallel.

**Theorem 11-B**

If two lines in a plane are cut by a transversal and the alternate exterior angles are congruent, then the lines are parallel.

**Theorem 11-C**

If two lines in a plane are cut by a transversal and the consecutive interior angles are supplementary, then the lines are parallel.

**Theorem 11-D**

If two lines in a plane are perpendicular to the same line, then the lines are parallel.

The distance from a point, which is not on a line, and a line is the length of a line segment that is perpendicular from the point to the line.

The distance between two parallel lines is the distance between one line and any point on the other line.

**Theorem 12-A  
Angle Sum  
Theorem**

The sum of the measures of the angles of a triangle is 180.

**Theorem 12-B  
Third Angle  
Theorem**

If two of the angles in one triangle are congruent to two of the angles in a second triangle, then the third angles of each triangle are congruent.

**Theorem 12-C  
Exterior Angle  
Theorem**

In a triangle, the measure of an exterior angle is equal to the sum of the measures of the two remote interior angles.

**Corollary 12-A-1**

The acute angles of a right triangle are complementary.

**Corollary 12-A-2**

There can be at most one right angle in triangle.

**Corollary 12-A-3**

There can be at most one obtuse angle in triangle.

**Corollary 12-A-4**

The measure of each angle in an equiangular triangle is 60.

**Definition of Congruent Triangles  
(CPCTC)**

Two triangles are congruent if and only if their corresponding parts are congruent.

**Postulate 12-A**

Any segment or angle is congruent to itself.  
(Reflexive Property)

**Postulate 13-A  
SSS Postulate**

If the sides of a triangle are congruent to the sides of a second triangle, then the triangles are congruent.

**SSS**

The three sides of one triangle must be congruent to the three sides of the other triangle.

**Postulate 13-B  
SAS Postulate**

If two sides and the included angle of a triangle are congruent to two sides and the included angle of a second triangle, then the triangles are congruent.

**SAS**

Two sides and the included angle of one triangle must be congruent to two sides and the included angle of the other triangle.

**Postulate 13-C  
ASA Postulate**

If two angles and the included side of a triangle are congruent to the two angles and included side of a second triangle, then the triangles are congruent.

**ASA**

Two angles and the included side of one triangle must be congruent to two angles and the included side of the other triangle.



**Theorem 13-A  
AAS Theorem**

**If two angles and a non-included side of a triangle are congruent to two angles and a non-included side of a second triangle, then the two triangles are congruent.**

**AAS**

Two angles and a non-included side of one triangle must be congruent to the corresponding two angles and side of the other triangle.

**Theorem 13-B  
Isosceles Triangle  
Theorem**

**If two sides of a triangle are congruent, then the angles that are opposite those sides are congruent.**

**Theorem 13-C**

**If two angles of a triangle are congruent, then the sides that are opposite those angles are congruent.**

**Corollary 13-B-1**

**A triangle is equilateral if and only if it is equiangular.**

**Corollary 13-B-2**

**Each angle of an equilateral triangle measures  $60^\circ$ .**

**Postulate 14-A  
HL Postulate**

**If there exists a correspondence between the vertices of two right triangles such that the hypotenuse and a leg of one triangle are congruent to the corresponding parts of the second triangle, then the two right triangles are congruent.**

**The shortest distance between two points is a straight line.**

**Postulate 14-B**

**A line segment is the shortest path between two points.**

**Theorem 14-A**

**A point on a perpendicular bisector of a segment is equidistant from the endpoints of the segment.**

**Theorem 14-B**

A point that is equidistant from the endpoints of a segment lies on the perpendicular bisector of the segment.

**Theorem 14-C**

A point on the bisector of an angle is equidistant from the sides of the angle.

**Theorem 14-D**

A point that is in the interior of an angle and is equidistant from the sides of the angle lies on the bisector of the angle.

**Comparison Property**

$a < b$ ,  $a = b$ , or  $a > b$ .

**Transitive Property**

1. If  $a < b$  and  $b < c$ , then  $a < c$ .
2. If  $a > b$  and  $b > c$ , then  $a > c$ .

**Addition Property**

1. If  $a > b$ , then  $a + c > b + c$ .
2. If  $a < b$ , then  $a + c < b + c$ .

**Subtraction Property**

1. If  $a > b$ , then  $a - c > b - c$ .
2. If  $a < b$ , then  $a - c < b - c$ .

**Multiplication Properties**

1. If  $c > 0$  and  $a < b$ , then  $ac < bc$ .
2. If  $c > 0$  and  $a > b$ , then  $ac > bc$ .
3. If  $c < 0$  and  $a < b$ , then  $ac > bc$ .
4. If  $c < 0$  and  $a > b$ , then  $ac < bc$ .

**Division Properties**

1. If  $c > 0$  and  $a < b$ , then  $\frac{a}{c} < \frac{b}{c}$ .
2. If  $c > 0$  and  $a > b$ , then  $\frac{a}{c} > \frac{b}{c}$ .
3. If  $c < 0$  and  $a < b$ , then  $\frac{a}{c} > \frac{b}{c}$ .
4. If  $c < 0$  and  $a > b$ , then  $\frac{a}{c} < \frac{b}{c}$ .

**Theorem 15-A  
Exterior Angle  
Inequality Theorem**

If an angle is an exterior angle of a triangle, then its measure is greater than the measure of either of its remote interior angles.

**Theorem 15-B**

If a side of a triangle is longer than another side, then the measure of the angle opposite the longer side is greater than the measure of the angle opposite the shorter side.

**Theorem 15-C**

In a triangle, if the measure of an angle is greater than the measure of a second angle, then the side that is opposite the larger angle is longer than the side opposite the smaller angle.

**Theorem 15-D**

The shortest segment from a point to a line is a perpendicular line segment between the point and the line.

**Theorem 15-E  
Triangle Inequality  
Theorem**

The sum of the lengths of any two sides of a triangle is greater than the length of the third side.

**Theorem 15-F  
SAS Inequality  
(Hinge Theorem)**

If two sides of a triangle are congruent to two sides of a second triangle, and if the included angle of the first triangle is greater than the included angle in the second triangle, then the third side of the first triangle is longer than the third side of the second triangle.

**Theorem 15-G  
SSS Inequality**

If two sides of a triangle are congruent to two sides of a second triangle, and if the third side in the first triangle is longer than the third side in the second triangle, then the included angle between the congruent sides in the first triangle is greater than the included angle between the congruent sides in the second triangle.

**Theorem 16-A**

The opposite sides of a parallelogram are congruent.

**Theorem 16-B**

The opposite angles of a parallelogram are congruent.

**Theorem 16-C**

The consecutive pairs of angles of a parallelogram are supplementary.

**Theorem 16-D**

**The diagonals of a parallelogram bisect each other.**

**Theorem 16-E**

**Either diagonal of a parallelogram separates the parallelogram into two congruent triangles.**

**Theorem 16-F**

**In a quadrilateral if both pairs of opposite sides are congruent, then the quadrilateral is a parallelogram.**

**Theorem 16-G**

**In a quadrilateral if both pairs of opposite angles are congruent, then the quadrilateral is a parallelogram.**

**Theorem 16-H**

**In a quadrilateral if its diagonals bisect each other, then the quadrilateral is a parallelogram.**

**Theorem 16-I**

**In a quadrilateral if one pair of opposite sides is both congruent and parallel, then the quadrilateral is a parallelogram.**

**Theorem 16-J**

**If a parallelogram is a rectangle, then its diagonals are congruent.**

**Theorem 16-K**

**If the diagonals of a parallelogram are congruent, then the parallelogram is a rectangle.**